



TITLE:

Geometric Characterizations of Standard Normal Distribution : Two Types of Differential Equations, Relationships with Square and Circle, and Their Similar Characterizations (Decision making theories under uncertainty and its applications : the extensions of mathematics for programming)

AUTHOR(S):

Nakanishi, Shingo; Ohnishi, Masamitsu

CITATION:

Nakanishi, Shingo ...[et al]. Geometric Characterizations of Standard Normal Distribution : Two Types of Differential Equations, Relationships with Square and Circle, and Their Similar Characterizations (Decision making theories under uncertainty ...

ISSUE DATE:

2018-07

URL:

<http://hdl.handle.net/2433/242116>

RIGHT:

Geometric Characterizations of Standard Normal Distribution

- Two Types of Differential Equations, Relationships with Square and Circle, and Their Similar Characterizations -

Shingo NAKANISHI

and

Masamitsu OHNISHI

Computing Center,
Osaka Institute of Technology

Graduate School of Economics and
Center for Mathematical Modeling and Data Science,
Osaka University

1. Introduction

We confirm various kinds of geometric characterizations of a standard normal distribution with Pearson's finding probabilistic point 0.612003 and its cumulative probability 0.2702678 on a standard normal distribution [1]. After Pearson proposed that, their numbers were studied by Kelley [2, 3], Mosteller [4], Cox [5] and many researchers [6-12]. Our studies [13-17] are also some of them. Sclove [6, 7] explained Cox's proposal to us as the clustering of normal distribution. Nakamori *et. al.* [8] also mentioned that Cox's proposal was one of the original papers of K-means Algorithms. Kelley's formulation, $\phi(\lambda) = 2\lambda\Phi(-\lambda)$, is also known as the 27 percent rule [9-12]. We found the other characterizations about 0.612003 from the parabolic curves of the cost of the repetitions game of coin tossing [13] and the square on a standard normal distribution with the probabilistic point 0.612003 [14-17].

In this paper, we confirm several characterizations based on this paper title. First, we show the geometric characterizations both Bernoulli differential equations [18] of inverse Mills ratios [19, 20] as the hazard functions and survival functions on standard normal distributions. At the same time, we find that Pearson's 0.612003 should be also an important role in their relations. Second, we clarify that variable coefficient second-order linear homogeneous differential equations [21] are related to a standard normal distribution. Especially, Sturm-Liouville form [22] with a probability density function of a standard normal distribution is shown concisely. Moreover, we illustrate that the value 0.612003 shows the curve which is the integral form of the cumulative distribution has the tangent lines on essential key points.

Furthermore, we also find the similar tendencies from European and Oriental historical cultures about the relations close to theology between circles and squares. Coincidentally, we get a practical approximated value 0.612004 from an equilateral triangle on the geometric characterization between circles, squares and a standard normal distribution instead of Pearson's finding true value 0.612003.




Figure 1 Concepts of integral forms of various kinds of cumulative distribution functions and their probability density functions. (a) Triangular distribution, (b) Uniform distribution, (c) Exponential distribution and (d) Standard normal distribution.

2. Our Concepts of This Study

Prof. Sandoh who was one of two referees of the first author's doctoral thesis [14] advised us to reconfirm the calculations practically about our previous research [13, 14] such as a general integral form [23]. We reconsider that as the following equation

$$E(U|U < -\lambda) = \frac{\int_{-\infty}^{-\lambda} u\phi(u)du}{\int_{-\infty}^{-\lambda} \phi(u)du} = -\frac{\phi(\lambda)}{\Phi(-\lambda)} = -2\lambda \quad (1)$$

as well as using the regression analysis [13, 14]. $\phi(u)$ is a probability density function of a standard normal distribution on

its probabilistic point u . $\Phi(-\lambda)$ is a calculated value of a cumulative distribution function from $-\infty$ to $-\lambda$ on u . λ that is about 0.612003 should be an important role of this study on u . $\phi(u)/\Phi(-u)$ is an inverse Mills ratio [19, 20]. Prof. Oya who was the other referee recommended us to investigate the tendencies between squares and normal distributions about our previous research [14]. We think of two precious advices as the concepts such as the integral forms of cumulative distribution functions like an idea by Nagasaka, Tsukamasa and Hashimoto [24] in Figure 1. That is, several purple 1-dot lines are shown in Figure 1. In this study, we clarify that the curve of an integral form of a cumulative distribution function of standard normal distribution has a principal role such as two types of differential equations and passes through several crucial points on their equations in this paper. By the way, Kelley also proposed the 27 percent rule formulation about $\lambda = 0.612003$ as follows

$$\frac{\phi(\lambda)}{\Phi(-\lambda)} = 2\lambda \text{ from } \frac{d\phi(u)}{du} = -u\phi(u) \text{ and } \frac{d\Phi(-u)}{du} = -\phi(u). \quad (2)$$

At this time, we show that the fact, $\phi(\lambda)$ is equal to $2\lambda\Phi(-\lambda)$ from Equations (1) and (2), brings us the equilibrium points in Figure 2. That is to say, a positive return is equal to its standard deviation as maximal loss [16, 17] and a negative return is equal to that as maximal profit [13-15, 17] under the condition that has the expected value as λ and its standard deviation 1.0 in Figure 2. From Figure 2, we confirm that the square whose length is 2λ is an expected value of a truncated normal distribution [25].




Figure 2 Equilibrium points both positive or negative returns and their risks under the condition on a standard normal distribution ($\lambda = 0.612003$, $t = 1.0$, Original reference [17]).

Especially, our concept about a negative return is similar to the meaningful insists of both Prof. Watanabe's website [26] and Prof. Tanioka's textbook [27] as the fee of gambling. On the other hand, the other about a positive return is similar to the general thinking about that a capital return is greater than its growth return historically such as Piketty's proposal [28].

3. Bernoulli Differential Equations for Inverse Mills Ratio

In section 2, we confirm that the equilibrium points on a standard normal distribution are shown based on $\lambda = 0.612003$. According to Isa's research [20], we rethink of the following formulations as Bernoulli differential equation with a standard normal distribution when we consider the tendencies of an inverse Mills ratio and a hazard function as the special cases on λ . The equation such as Bernoulli differential equation with a standard normal distribution is also mentioned in Prof. Kijima's special lecture [18]. However, since we are interested in the geometric characterizations of that, we propose two types of homogeneous Bernoulli differential equations. That is




Figure 3 Bernoulli differential equation of the ratios of probability density function of standard normal distribution out of its cumulative distribution function with thinking of its truncated normal distribution on the probabilistic point $\lambda = 0.612003$.

$$\frac{dg(u)}{du} + ug(u) - g(u)^2 = 0, \quad g(u) = \frac{\phi(u)}{C + \Phi(-u)}, \quad (3)$$

$$\frac{dg(u)}{du} + ug(u) + g(u)^2 = 0, \quad g(u) = \frac{\phi(u)}{C + \Phi(u)}. \quad (4)$$

Based on the ideas of asymptotic lines by Gordon [19] and Isa [20], we illustrate the Bernoulli differential equations as the geometric curves in the upper chart of Figure 3 with $\lambda = 0.612003$. At this time, a constant C is changed by k from 0.0 to 3.0 by 0.1 to illustrate those curves clearly. Especially, we confirm that their tendencies are shown as the special types with $\lambda = 0.612003$ when $k = 0.0$ or 1.0. Moreover, 0.30263084 that is also a probabilistic point with the same values of both its cumulative distribution function and probability density function is another important point because the inverse Mills ratio is equal to 1.0. Under the condition of $u = 0$, we find the inverse Mills ratio is $2\phi(0)(= 2/\sqrt{2\pi})$ as 2 times of $\phi(0)$. The lower chart also illustrates the geometric tendencies about Equations (3) and (4) divided by $u = 0$ symmetrically.

4. Sturm-Liouville Differential Equations for a Standard Normal Distribution

In section 2, we mentioned our research concepts as the integral forms of a cumulative distribution of a standard normal distribution. We searched for their several characterizations [29, 30] and their differential equations about normal distribution [30, 31]. From our investigations, we propose the variable coefficient second-order linear homogeneous differential equation in Figure 4.

Sturm-Liouville differential equation: $\phi(u)(h_2'(u) - uh_2'(u) - h_2(u)) = 0$

Sturm-Liouville form: $(\phi(u)h_2'(u))' - (\phi(u)h_2(u)) = 0$

Variable coefficient second-order linear homogeneous differential equation for

a standard normal distribution: $\frac{d^2 h_2(u)}{du^2} - u \frac{dh_2(u)}{du} - h_2(u) = 0,$

$h_2(u)$

$f(u)$

$\phi(u)$

$\psi(u)$


ϕ_2

$$\frac{dh_2(u)}{du} = -\Phi(-u) \quad \frac{d^2 h_2(u)}{du^2} = \phi(u)$$

Equilibrium point
between a banker, winners
and losers: $\phi(\lambda) = 2\lambda\Phi(-\lambda)$

$$\lambda = 0.612003$$

Probability density
function of a truncated
normal distribution: $\psi(u) = \frac{\phi(u)}{\Phi(-\lambda)}$



$$\text{Tangential equation: } f(u) = \begin{cases} -(1-\Phi(-\lambda))u + 2\lambda\Phi(-\lambda) & (-2 \leq u \leq 0) \\ -\Phi(-\lambda)u + 2\lambda\Phi(-\lambda) & (0 \leq u \leq 2\lambda) \end{cases}$$

Probability density function of a standard normal distribution: $\phi(u)$

Cumulative distribution function of a standard normal distribution: $\Phi(-u)$

Equilibrium point

of an inverse

$$\text{Mills ratio: } \frac{\phi(\lambda)}{\Phi(-\lambda)} = 2\lambda$$

$$\text{Intercept form of a linear equation for winners: } -\frac{1}{\lambda}u + \frac{1}{\lambda\Phi(-\lambda)}\phi_2 = 1$$

$$\text{Intercept form of a linear equation for losers: } -\frac{1}{\lambda}u + \frac{1}{\lambda(1+\Phi(-\lambda))}\phi_2 = 1$$

$$\text{Intercept form of a linear equation for a banker: } -\frac{1}{\lambda}u + \frac{1}{\lambda}\phi_2 = 1$$

Utility function for winners:

$$U_w(t) = (\phi(\lambda) - \lambda\Phi(-\lambda))\sqrt{t} = \lambda\Phi(-\lambda)\sqrt{t}$$

Figure 4 Geometric relationships between Sturm-Liouville differential equation and standard normal distribution with a circle, squares, meaningful rectangles, their diagonals, and special tangent lines under $\lambda = 0.612003$.

That is

$$\frac{d^2 h_2(u)}{du^2} - u \frac{dh_2(u)}{du} - h_2(u) = 0. \quad (5)$$

The way of an expression of $h_2(u)$ in Equation (5) is based on the descriptions of our previous research [14]. The formulations of $h_2(u)$ are shown as the following equation

$$h_2(u) = A(\phi(u) - u\Phi(-u)) + B(\phi(u) + u\Phi(u)). \quad (6)$$

If we consider the conditions $dh_2(u)/du = 1/\sqrt{2\pi}$ and $h_2(u) = 1/2$ in Equation (5), we get the answer with $A = 1.0$ and $B = 0.0$ in Equation (6). Therefore, we also confirm that the special curve in Figure 4 passes through several fundamental points with $\lambda\Phi(-\lambda)$ and $\lambda(1 + \Phi(-\lambda))$ [17]. At the same time, we find their points also have the crucial tangent lines shown in Figure 4. Moreover, we illustrate a true circle with a square and a standard normal distribution in Figure 4 [17]. The center of this circle is an intersection of meaningful rectangle diagonals. Without passing through the intersection, we find the other important line to distinguish into the separated squares by the proportion $\Phi(-\lambda): 1 - \Phi(-\lambda)$.

We reconfirm the geometric characterizations about $\lambda = 0.612003$ with considering a square, a circle and a normal distribution. And their tendencies have a special differential equation as the following Sturm-Liouville form [22]

$$(\phi(u)h_2'(u))' + (\phi(u)h_2(u)) = 0. \quad (7)$$

At this time, to investigate the tendencies of the combinations between A and B in Equation (6), A and B are changed by the values from 0.0 to 1.0 by 0.25 respectively in Figure 5.




Figure 5 Various geometric characterizations of Sturm-Liouville differential equations with standard normal distribution.

5. Square and Circle on Standard Normal Distribution and Approximated Value about Polygons

Although we confirm that there is a relationship between a circle, a square and a normal distribution, we cannot get a correct value λ as well as drawing an equilateral triangle and a regular hexagon from their characterizations which are similar to the problem of squaring the circle [32, 33]. Instead of that, we can get the practical approximated Pearson's finding value. That is

$$\lambda \cong \frac{1 + \sqrt{3}}{1 + 2\sqrt{3}} = 0.612004. \quad (8)$$

This approximated value is estimated by drawing equilateral triangles and regular hexagons shown in Figure 6.

By the way, there are a lot of thinking about circles and squares for a long time all over the world religiously and historically [32-39]. We also show one of their tendencies with a standard normal distribution based on $\lambda = 0.612003$. Coincidentally,

を記します。

References

- [1] Pearson, K., "On the probable errors of frequency constants", *Biometrika*, **13**, 113-32, (1920).
- [2] Kelley, T. L., "Footnote 11 to Jensen, MB, Objective differentiation between three groups in education (teachers, research workers, and administrators)", *Genetic psychology monographs*, **3**(5), 361, (1928).
- [3] Kelley, T. L., "The selection of upper and lower groups for the validation of test items", *Journal of Educational Psychology*, **30**(1), 17-24 (1939).
- [4] Mosteller, F., "On Some Useful Inefficient Statistics", *The Annals of Mathematical Statistics*, **17**, 377-408, (1946).
- [5] Cox, D. R., "Note on grouping", *Journal of the American Statistical Association*, **52**(280), 543-547, (1957).
- [6] Sclove, S. L., "A Course on Statistics for Finance", Chapman and Hall/CRC, (2012).
- [7] Johari, S. and Sclove, S. L., "Partitioning a distribution", *Communications in Statistics*, **A5**, 133-147, (1976).
- [8] Mar, S. O., Huynh, V.-N. and Nakamori, Y., "An alternative extension of the k-means algorithm for clustering categorical data", *International Journal of Applied Mathematics and Computer Science*, **14**(2), 241-248, (2004).
- [9] Cureton, E. E., "The upper and lower twenty-seven per cent rule", *Psychometrika*, **22**(3), 293-296, (1957).
- [10] Ross, J. and Weitzman, R. A., "The twenty-seven per cent rule", *The Annals of Mathematical Statistics*, 214-221, (1964).
- [11] 赤根敦, 伊藤圭, 林篤裕, 椎名久美子, 大澤公一, 柳井晴夫, 田栗正章, "識別指数による総合試験問題の項目分析", 大学入試センター研究紀要, **35**, 19-47, (2006).
- [12] 豊田秀樹, 秋山隆, 岩間徳兼, "項目特性図における情報量規準を用いた群数の選択法", 教育心理学研究, **62**(3), 209-225, (2014).
- [13] Nakanishi, S., "Maximization for sum total expected gains by winners of repetition game of coin toss considering the charge based on its number trial", *Transactions of the Operations Research Society of Japan*, **55**, 1-26, (2012) in Japanese.
- [14] Nakanishi, S., "Uncertainties of active management in the stock portfolios in consideration of the fee", *Doctoral Thesis, Graduate School of Economics, Osaka University*, (17777), (2015) in Japanese.
- [15] 中西真悟, "27 パーセントルールと逆ミルズ比に基づく手数料を考慮した繰り返しコイン投げゲームの勝者の最大賞金と正規分布の関係", 日本 OR 学会 2016 年春季研究発表会アブストラクト集, 115-116, (2016).
- [16] Nakanishi, S., "Equilibrium point on standard normal distribution between expected growth return and its risk", *Abstract of 28th European Conference on Operational Research (the EURO XXVIII)*, Poznan, Poland, (2016).
- [17] 中西真悟, 大西匡光, "27 パーセントルールと逆ミルズ比を用いたリスクと正負のリターン統計学的特性と幾何学的特性", 日本 OR 学会 2017 年秋季研究発表会アブストラクト集, 181-182, (2017).
- [18] 木島正明, "招待講演資料: A Model of Price Impact Function", 日本 OR 学会 4 部会合同研究会 ～確率モデルの新展開～, (2017).
- [19] Gordon, R. D., "Values of Mills' Ratio of Area to Bounding Ordinate and of the Normal Probability Integral for Large Values of the Argument", *The Annals of Mathematical Statistics*, **12**(3), 364-366, (1941).
- [20] 伊佐勝秀, "ミルズ比, 逆ミルズ比及び正規ハザード率 — その諸性質と経済学における応用に関する部分的サーベイ —", 西南学院大学経済学論集, 西南大学学術研究所, **45**(4), 131-150, (2011).
- [21] Tenenbaum, M. and Pollard, H., "Ordinary Differential Equations", Dover Publications, (1985).
- [22] Al-Gwaiz, M., "Sturm-Liouville Theory and its Applications", Springer, (2007).
- [23] Hosseiniour, A. and Sandoh, H., "Optimal business hours of the newsvendor problem for retailers", *International Transactions in Operational Research*, **20**, 823-836, (2013).
- [24] 長坂一徳, 塚正勉, 橋本文雄, "工具の損耗を考慮した最適工具交換時期の設定", 精密機械, **51**(6), 1207-1211, (1985).
- [25] Ahsanullah, M., Kibria, B.M.G. and Shakil, M., "Normal and Student's t Distributions and Their Applications", Atlantis Press, (2014).
- [26] 渡辺隆裕, "賭けの科学", <http://www.nabenavi.net/>, (Access month: March, 2010).
- [27] 谷岡一郎, "ツキの法則-「賭け方」と「勝敗」の科学", PHP 研究所, (1997).
- [28] Piketty, T., (山形浩生, 守岡桜, 森本正史 翻訳), "21 世紀の資本", みすず書房, (2014).
- [29] 柴田義貞, "正規分布 特性と応用", 東京大学出版会, (1981).
- [30] Bryc, W., "The Normal Distribution: Characterizations with Applications", Springer New York, (1995).
- [31] 兼清泰明, "確率微分方程式とその応用", 森北出版, (2017).
- [32] Wikipedia, "円積問題", <https://ja.wikipedia.org/wiki/円積問題>, (Access month, November, 2017) in Japanese.
- [33] Wikipedia, "Squaring the Circle", https://en.wikipedia.org/wiki/Squaring_the_circle, (Access month, November, 2017).
- [34] 井田隆, "ダ・ヴィンチのウィトルウィウスの人体図と黄金比", <http://www.crl.nitech.ac.jp/~ida/education/VitruvianMan/index-j.html>, (Access month: July, 2017).
- [35] Ida, T., "Vitruvian Man by Leonardo da Vinci and the Golden Ratio", (Access month: July, 2017). <http://www.crl.nitech.ac.jp/~ida/education/VitruvianMan/index.html>, (Access month: July, 2017).
- [36] Wikipedia, "Mandala", <https://en.wikipedia.org/wiki/Mandala>, (Access month: July, 2017).
- [37] Wikipedia, "曼荼羅", <https://ja.wikipedia.org/wiki/曼荼羅>, (Access month: July, 2017) in Japanese.
- [38] Wikipedia, "正規分布", <https://ja.wikipedia.org/wiki/正規分布>, (Access month, July, 2017) in Japanese.
- [39] Wikipedia, "Normal distribution", https://en.wikipedia.org/wiki/Normal_distribution, (Access month, July, 2017).